arXiv:1409.3932v2 [math-ph] 10 Mar 2015

THE PADÉ INTERPOLATION METHOD APPLIED TO q-PAINLEVÉ EQUATIONS

HIDEHITO NAGAO

Abstract. We establish interpolation problems related to all the q-Painlevé equations of types from $E_7^{(1)}$ to $(A_2 + A_1)^{(1)}$. By solving those problems, we can derive the evolution equations, the scalar Lax pairs and the determinant formulae of special solutions for the corresponding q-Painlevé equations.

Contents

1. Introduction 1
1.1. The background of discrete Painlevé equations 1
1.2. The background of the Padé method 2
1.3. The purpose and the organization of this paper 2
2. Padé interpolation method 3
2.1. (a) Setting of the Padé interpolation problem 3
2.2. (b) Computation of contiguity relations 4
2.3. (c) Computation of the q-Painlevé equation 5
2.4. (d) Computation of the Lax pair 5
2.5. (e) Computation of special solutions 6
3. Main results 7
3.1. Case q-$E_7^{(1)}$ 7
3.2. Case q-$E_6^{(1)}$ 9
3.3. Case q-$D_5^{(1)}$ 11
3.4. Case q-$A_4^{(1)}$ 12
3.5. Case q-$(A_2 + A_1)^{(1)}$ 13
4. Conclusion 15
4.1. Summary 15
4.2. Problems 15
Acknowledgment 15
References 15

1. Introduction

1.1. The background of discrete Painlevé equations.

Discrete Painlevé equations are discrete equations which reduce to differential Painlevé equations by a suitable limiting process. As integrable systems, they are studied from various points of view (see [3] for example).

In Sakai's theory [19], the discrete Painlevé equations were classified on the basis of rational surfaces connected to extended affine Weyl groups. There exist three types of discrete Painlevé equations in the classification: elliptic difference (e-), multiplicative difference (q-) and additive difference (d-). The discrete Painlevé equations of q-difference type are classified as follows:

2010 *Mathematics Subject Classification.* 33D15, 34M55, 39A13, 41A21.
Key words and phrases. Padé method, Padé interpolation, q-Painlevé equation.

$$E_8^{(1)} \to E_7^{(1)} \to E_6^{(1)} \to \underset{(q\text{-}P_{\mathrm{VI}})}{D_5^{(1)}} \to \underset{(q\text{-}P_{\mathrm{V}})}{A_4^{(1)}} \to \underset{(q\text{-}P_{\mathrm{IV}},q\text{-}P_{\mathrm{III}})}{(A_2+A_1)^{(1)}} \to \underset{(q\text{-}P_{\mathrm{II}})}{(A_1+A_1')^{(1)}} \to \underset{(q\text{-}P_{\mathrm{I}})}{A_1^{(1)}} \to \mathcal{D}_6$$

$$\nearrow \mathbb{Z}$$

1.2. **The background of the Padé method.**

1.2.1. *What is the Padé method?*

The method, which we call the *Padé method* in this paper, is a method for giving the Painlevé equations, the scalar Lax pairs and the determinant formulae of special solutions simultaneously, by starting from suitable problems of Padé approximation/interpolation.

There exists a close connection between Painlevé equations and Padé approximation/interpolation. The Padé method has been presented by applying Padé approximation to differential Painlevé equations of types from P_{VI} to P_{IV} in [22].

The Padé method is closely related to the theory of orthogonal polynomials (e.g.,[1][12][26][28]). By both approaches, we can obtain Painlevé equations, Lax pairs and special solutions. (The theory of orthogonal polynomials is more general and the Padé method is simpler.)

1.2.2. *Previous works on the Padé method.*

Concerning all the q-Painlevé equations of types from $E_7^{(1)}$ to $(A_2+A_1)^{(1)}$ in the above classification, the q-Painlevé equations, the scalar Lax pairs and the special solutions have been already derived by various methods. References are written in the following Table (1.1):

(1.1)

	$q\text{-}E_7^{(1)}$	$q\text{-}E_6^{(1)}$	$q\text{-}D_5^{(1)}$	$q\text{-}A_4^{(1)}$	$q\text{-}(A_2+A_1)^{(1)}$
q-Painlevé	[4]	[17]	[8]	[11]	[10][11][19]
Lax pair	[23]	[21]	[8]	[14]	[14]
Special solutions	[13]	[6]	[18]	[5]	[15]

Here, amongst the references in the Table (1.1), only [6] is directly related to the Padé method.

The Padé method for discrete Painlevé equations has been applied to the following cases:

(1.2)

	$e\text{-}E_8^{(1)}$	$q\text{-}E_8^{(1)}$	$q\text{-}E_6^{(1)}$	$q\text{-}D_5^{(1)}$
	[16]	[24]	[6]	[6][25]
Grid	elliptic	q-quadric	q	differential, q

Here, the case $q\text{-}D_5^{(1)}$ is studied as both the q-grid and the differential grid (i.e., Padé approximation).

1.3. **The purpose and the organization of this paper.**

The purpose of this paper is to apply the Padé method to all the q-Painlevé equations of types from $q\text{-}E_7^{(1)}$ to $q\text{-}(A_2+A_1)^{(1)}$. Namely, by the Padé method, we derive the evolution equations, the scalar Lax pairs and the determinant formulae of special solutions for the corresponding q-Painlevé equations. As the main results, the following items are presented for each type.

(a) Setting of the Padé interpolation problem,
(b) Contiguity relations,
(c) The q-Painlevé equation,
(d) The Lax pair,
(e) Special solutions.

This paper is organized as follows: In Section 2, we will explain the Padé interpolation method applied to the q-Painlevé equations. Namely, we will explain the methods for deriving items (a)–(e). In Section 3, we will present these main results for all the q-Painlevé equations of types from $E_7^{(1)}$ to $(A_2+A_1)^{(1)}$. In Section 4, we will give a summary and discuss some future problems.

2. Padé interpolation method

In this section, we will explain the methods for deriving the items (a)–(e) in the main results given in Section 3.

2.1. **(a) Setting of the Padé interpolation problem.**

• In this section, we will consider the following interpolation problem (of the q-grid):

For a given function $Y(x)$, we look for functions $P_m(x)$ and $Q_n(x)$ which are polynomials of degree m and n, satisfying the interpolation condition

$$Y(q^s)=P_m(q^s)/Q_n(q^s) \quad (s=0,1,\dots,m+n). \tag{2.1}$$

We call this problem the *Padé interpolation problem (of the q-grid)*. Then, the function $Y(x)$ is called the *interpolated function*. Correspondingly, the polynomials $P_m(x)$ and $Q_n(x)$ are called *interpolating functions* respectively.

The common normalization factor of the polynomials $P_m(x)$ and $Q_n(x)$ is not determined by the condition (2.1). However, this normalization factor is not essential for our arguments, i.e., the main results in Section 3 (see Remark 2). The explicit expressions of $P_m(x)$ and $Q_n(x)$, which will be used in the computations for the item (e) above, were essentially given in [7] (see item (e) below).

• In this section, we will establish the interpolation problems (2.1) by specifying the interpolated functions $Y(x)$ and the interpolated sequences $Y_s=Y(q^s)$ as follows:

(2.2)

	q-$E_7^{(1)}$	q-$E_6^{(1)}$	q-$D_5^{(1)}$	q-$A_4^{(1)}$	q-$(A_2+A_1)^{(1)}$
$Y(x)$	$\prod_{i=1}^{3}\frac{(a_ix,b_i;q)_\infty}{(a_i,b_ix;q)_\infty}$ $\frac{a_1a_2a_3q^m}{b_1b_2b_3q^n}=1$	$\prod_{i=1}^{2}\frac{(a_ix,b_i;q)_\infty}{(a_i,b_ix;q)_\infty}$	$c^{\log_q x}\frac{(a_1x,b_1;q)_\infty}{(a_1,b_1x;q)_\infty}$	$c^{\log_q x}\frac{(b_1;q)_\infty}{(b_1x;q)_\infty}$	$(d\sqrt{x/q})^{\log_q x}$
Y_s	$\prod_{i=1}^{3}\frac{(b_i;q)_s}{(a_i;q)_s}$	$\prod_{i=1}^{2}\frac{(b_i;q)_s}{(a_i;q)_s}$	$c^s\frac{(b_1;q)_s}{(a_1;q)_s}$	$c^s(b_1;q)_s$	$q^{\binom{s}{2}}d^s$
q-HGF	${}_4\varphi_3$	${}_3\varphi_2$	${}_2\varphi_1$	${}_2\varphi_1$	${}_1\varphi_1$

where $\frac{a_1a_2a_3q^m}{b_1b_2b_3q^n}=1$ is a constraint for the parameters in the case q-$E_7^{(1)}$, and the q-shifted factorials are defined by

$$(a_1,a_2,\dots,a_i;q)_j=\prod_{k=0}^{j-1}(1-a_1q^k)(1-a_2q^k)\dots(1-a_iq^k), \tag{2.3}$$

and the q-HGF (the q-hypergeometric functions [2]) is defined by

$${}_k\varphi_l\left(\begin{matrix}a_1,&\dots,&a_k\\ b_1,&\dots,&b_l\end{matrix};q,x\right)=\sum_{s=0}^{\infty}\frac{(a_1,\dots,a_k;q)_s}{(b_1,\dots,b_l,q;q)_s}\left[(-1)^sq^{\binom{s}{2}}\right]^{1+l-k}x^s, \tag{2.4}$$

with $\binom{s}{2}=s(s-1)/2$.

Remark 1. On the choice of Y_s and $Y(x)$
In Table (2.2), the expressions for the interpolated sequences Y_s are closely related to those of q-HGF ${}_r\varphi_s$. In fact, we chose the Y_s by comparing the coefficient of x^s in ${}_r\varphi_s$ (2.4) and the expression for $Y_s(q^{-(m+n)};q)_s/(q;q)_s$ in (2.15). Then, we also chose the interpolated functions $Y(x)$, which are equal to the Y_s at $x=q^s$. □

• In this section, we will consider yet another Padé problem where some parameters a_i, b_i, c, d, m and n in $Y(x)$ are shifted. The parameter shift operators T are given as follows:

(2.5)

	Parameters		
q-$E_7^{(1)}$	$(a_1,a_2,a_3,b_1,b_2,b_3,m,n)$	$\mapsto$	$(qa_1,a_2,qa_3,b_1,b_2,qb_3,m-1,n)$
q-$E_6^{(1)}$	(a_1,a_2,b_1,b_2,m,n)	$\mapsto$	$(qa_1,a_2,b_1,b_2,m-1,n)$
q-$D_5^{(1)}$	(a_1,b_1,c,m,n)	$\mapsto$	$(qa_1,b_1,c,m-1,n)$
q-$A_4^{(1)}$	(b_1,c,m,n)	$\mapsto$	$(b_1,c,m-1,n)$
q-$(A_2+A_1)^{(1)}$	(d,m,n)	$\mapsto$	$(d,m-1,n)$

Here, the operators T are called the *time evolutions*, because they specify the directions of the time evolutions for q-Painlevé equations.

2.2. **(b) Computation of contiguity relations.**

• We will consider the following pair of linear q-difference equations $L_2(x)$ and $L_3(x)$ for unknown function $y(x)$, which are the main object in our study.

$$L_2(x): \begin{vmatrix} y(x) & y(qx) & \overline{y}(x) \\ P_m(x) & P_m(qx) & \overline{P}_m(x) \\ Y(x)Q_n(x) & Y(qx)Q_n(qx) & \overline{Y}(x)\overline{Q}_n(x) \end{vmatrix} = 0,$$

(2.6)
$$L_3(x): \begin{vmatrix} y(x) & \overline{y}(x) & \overline{y}(x/q) \\ P_m(x) & \overline{P}_m(x) & \overline{P}_m(x/q) \\ Y(x)Q_n(x) & \overline{Y}(x)\overline{Q}_n(x) & \overline{Y}(x/q)\overline{Q}_n(x/q) \end{vmatrix} = 0.$$

Then, the pair of q-difference equations (2.6) is called the *contiguity relations*. Here, $\overline{F}$ and $\underline{F}$ denote $T(F)$ and $T^{-1}(F)$ respectively. T is the shift operator acting on parameters given in Table (2.5).

• We will show the method of computation of the contiguity relations $L_2(x)$ and $L_3(x)$.
Set $\mathbf{y}(x)=\begin{bmatrix} P_m(x) \\ Y(x)Q_n(x) \end{bmatrix}$ and define Casorati determinants $D_i(x)$ by

(2.7)
$$D_1(x)=\det[\mathbf{y}(x),\mathbf{y}(qx)],\quad D_2(x)=\det[\mathbf{y}(x),\overline{\mathbf{y}}(x)],\quad D_3(x)=\det[\mathbf{y}(qx),\overline{\mathbf{y}}(x)].$$

Then, the contiguity relations (2.6) can be rewritten as follows:

(2.8)
$$\begin{aligned} &L_2(x): D_1(x)\overline{y}(x)-D_2(x)y(qx)+D_3(x)y(x)=0,\\ &L_3(x): \overline{D}_1(x/q)y(x)+D_3(x/q)\overline{y}(x)-D_2(x)\overline{y}(x/q)=0. \end{aligned}$$

We define basic quantities $G(x), K(x)$ and $H(x)$ [e.g., (3.4), (3.17)] by

(2.9)
$$G(x)=Y(qx)/Y(x),\quad K(x)=\overline{Y}(x)/Y(x),\quad H(x)=\mathrm{L.C.M}(G_{\mathrm{den}}(x),K_{\mathrm{den}}(x)),$$

where $G_{\mathrm{den}}(x)$ and $G_{\mathrm{num}}(x)$ are defined as the polynomials of the denominator and the numerator of $G(x)$ respectively, and $K_{\mathrm{den}}(x)$ and $K_{\mathrm{num}}(x)$ are similarly defined. Substituting these quantities into the

equations (2.7), we obtain the following determinants:

$$
\begin{aligned}
D_1(x) &= \frac{Y(x)}{G_{\mathrm{den}}(x)}\{G_{\mathrm{num}}(x)P_m(x)Q_n(qx) - G_{\mathrm{den}}(x)P_m(qx)Q_n(x)\},\\
D_2(x) &= \frac{Y(x)}{K_{\mathrm{den}}(x)}\{K_{\mathrm{num}}(x)P_m(x)\overline{Q}_n(x) - K_{\mathrm{den}}(x)\overline{P}_m(x)Q_n(x)\},\\
D_3(x) &= \frac{Y(x)}{H(x)}\{\frac{H(x)}{K_{\mathrm{den}}(x)}K_{\mathrm{num}}(x)P_m(qx)\overline{Q}_n(x) - \frac{H(x)}{G_{\mathrm{den}}(x)}G_{\mathrm{num}}(x)\overline{P}_m(x)Q_n(qx)\}.
\end{aligned}
\tag{2.10}
$$

Using the interpolation condition (2.1) and the form of the basic quantities [e.g., (3.4), (3.17)], we can investigate positions of zeros and degrees of the polynomials within braces $\{\ \ \}$ of the equations (2.10). Then, we can simply compute the determinants $D_i(x)$ [e.g., (3.5), (3.18)] except for some factors such as $1 - fx$ in $D_1(x)$ and $1 - x/g$ in $D_3(x)$, where f and g, etc, are constants with respect to x. In this way, we obtain the contiguity relations $L_2(x)$ and $L_3(x)$ [e.g., (3.6), (3.19)].

Remark 2. On the gauge invariance of C_0C_1
When the common normalization factor of $P_m(x)$ and $Q_n(x)$ is changed, an x-independent gauge transformation of $y(x)$ is induced in $L_2(x)$ and $L_3(x)$. Under the x-independent gauge transformation of $y(x)$: $y(x) \mapsto Gy(x)$, the coefficients of $\overline{y}(x), y(x/q), y(x)$ and $y(x), \overline{y}(x), \overline{y}(x/q)$ in (2.8) change as follows:

$$
\begin{aligned}
&(D_1(x) : D_2(x) : D_3(x)) \mapsto (\overline{G}D_1(x)/G : D_2(x) : D_3(x))\\
&(\overline{D}_1(x/q) : D_3(x/q) : D_2(x)) \mapsto (G\overline{D}_1(x/q)/\overline{G} : D_3(x/q) : D_2(x)).
\end{aligned}
\tag{2.11}
$$

The coefficients C_0 and C_1 in $L_2(x)$ and $L_3(x)$ [e.g., (3.6), (3.19)] are defined as the normalization factors of the coefficients of $\overline{y}(x)$ and $y(x)$ respectively. Then, C_0 and C_1 change under the gauge transformation, but the product C_0C_1 is a gauge invariant quantity. Moreover, C_0 and C_1 do not appear in the final form of the q-Painlevé equations. □

2.3. **(c) Computation of the q-Painlevé equation.**

We can derive the q-Painlevé equation from the compatibility condition of the contiguity relations $L_2(x)$ and $L_3(x)$ [e.g., (3.6), (3.19)]. Computing the compatibility condition, we determine three quantities $\underline{g}$, $\overline{f}$ and C_0C_1. Expressions for $\underline{g}$ and $\overline{f}$ are obtained in terms of f and g. An expression for C_0C_1 is obtained in terms of f, g and $\overline{f}$ (and hence in terms of f and g).

The first and the second expressions are the q-Painlevé equation [e.g., (3.8), (3.20)]. The third expression is a constraint for the product C_0C_1 [e.g., (3.10), (3.22)].

Remark 3. On two meanings of f, g, m and n
We use the variables f and g with two different meanings. The first meaning is the f and g which are explicitly determined in terms of parameters a_i, b_i, m and n by the Padé problem. The second meaning is the f and g which are unknown functions in the q-Painlevé equation. In items (c) and (d), we consider f and g in the second meaning.

Similarly, we use the parameters m and n with two meanings. In the first meaning, m and n are integers. In the second meaning, m and n are generic complex parameters. In items (c) and (d), we consider m and n in the second meaning. Then, the result of the compatibility of $L_2(x)$ and $L_3(x)$ also holds with respect to the second meaning. □

2.4. **(d) Computation of the Lax pair.**

• We will consider the following pair of linear q-difference equations for the unknown function $y(x)$:

$$
\begin{aligned}
&L_1(x) : A_1(x)y(x/q) + A_2(x)y(x) + A_3(x)y(qx) = 0,\\
&L_2(x) : A_4(x)\overline{y}(x) + A_5(x)y(x) + A_6(x)y(qx) = 0,
\end{aligned}
\tag{2.12}
$$

such that their compatibility condition gives a q-Painlevé equation. Then, the pair of q-difference equations is called the "scalar Lax pair".

• We will show the method of computation of the scalar Lax pair (2.12).
The Lax pair $L_1(x)$ and $L_2(x)$, which satisfies the compatibility condition, is derived using the results of items (a)–(c) as follows: The $L_2(x)$ equation in item (d) [e.g., (3.11), (3.23)] is the same as the $L_2(x)$ in the item (b). We can obtain the Lax equation $L_1(x)$ as follows: First, combining the contiguity relations $L_2(x)$ and $L_3(x)$ [e.g., (3.11), (3.23)], one obtains an equation between the three terms $y(qx), y(x)$ and $y(x/q)$ (See the figure below), whose coefficient functions depend on the variables $f, g, \overline{f}, C_0$ and C_1. However, the variables C_0 and C_1 appear through the product C_0C_1. Therefore, expressing $\overline{f}$ and C_0C_1 in terms of f and g only, one obtains the Lax equation $L_1(x)$ [e.g., (3.11), (3.23)].



2.5. **(e) Computation of special solutions.**

By construction, expressions for f and g as in the first meaning in Remark 3 give a special solution for the q-Painlevé equation. We will present how to compute determinant formulae of the special solutions.

• We will derive formulae (2.15) which are convenient for computing the special solutions f and g. The Cauchy–Jacobi formulae (2.14) are essentially presented in [7]. For a given sequence Y_s, the polynomials $P_m(x)$ and $Q_n(x)$ of degree m and n for an interpolation problem

$$Y_s = P_m(x_s)/Q_n(x_s) \quad (s = 0, 1, \dots, m+n) \tag{2.13}$$

are given by the following determinant expressions:

$$P_m(x) = F(x) \det\Big[\sum_{s=0}^{m+n} u_s \frac{x_s^{i+j}}{x - x_s} \Big]_{i,j=0}^{n}, \quad Q_n(x) = \det\Big[\sum_{s=0}^{m+n} u_s x_s^{i+j} (x - x_s) \Big]_{i,j=0}^{n-1}, \tag{2.14}$$

where $u_s = Y_s/F'(x_s)$ and $F(x) = \prod_{i=0}^{m+n}(x - x_i)$.

In the q-grid case of problem (2.13) (i.e., the case of problem (2.1)), the formulae (2.14) take the following form:

$$\begin{aligned} P_m(x) &= \frac{F(x)}{(q;q)_{m+n}^{n+1}} \det\Big[\sum_{s=0}^{m+n} Y_s \frac{(q^{-(m+n)};q)_s}{(q;q)_s} \frac{q^{s(i+j+1)}}{x - q^s} \Big]_{i,j=0}^{n}, \\ Q_n(x) &= \frac{1}{(q;q)_{m+n}^{n}} \det\Big[\sum_{s=0}^{m+n} Y_s \frac{(q^{-(m+n)};q)_s}{(q;q)_s} q^{s(i+j+1)} (x - q^s) \Big]_{i,j=0}^{n-1}. \end{aligned} \tag{2.15}$$

In the derivation of (2.15), we have used the following relations:

$$\begin{aligned} F'(x_s) =& (x_s - x_0) \dots (x_s - x_{s-1})(x_s - x_{s+1}) \dots (x_s - x_{m+n}) \\ =& (-1)^s q^{(s-1)s/2} (q;q)_s q^{s(m+n-s)} (q;q)_{m+n-s} \\ =& (q;q)_s (q;q)_{m+n} / q^s (q^{-(m+n)};q)_s. \end{aligned} \tag{2.16}$$

Moreover, substituting the values of Y_s (2.2) and $F'(x_s)$ (2.16) into the formulae (2.14), one obtains the determinant formulae (2.15).

• We will show the method of computation of the special solutions f and g.
The expressions for the f and g can be derived by comparing $D_i(x)$ in (2.10) and $D_i(x)$ [e.g., (3.5), (3.18)] in item (b) as the identity with respect to the variable x and applying the formulae (2.15).

For example, the computation for the case q-$E_7^{(1)}$ is as follows: Substituting $x = 1/a_i$ into $D_1(x)$ in (2.10) and $D_1(x)$ in (3.5), we obtain an expression for the f in the first equation of (3.12) by comparing the two expressions for $D_1(x)$ and applying the formulae (2.15). Similarly, substituting $x = 1/b_i$ into the third equation of (2.10) and $D_3(x)$ in (3.5), we obtain an expression for the g in the second equation of (3.12) by comparing the two expressions for $D_3(x)$ and applying the formulae (2.15).

3. Main results

In this subsection, we will present the results obtained through the method, which is explained for each case from q-$E_7^{(1)}$ to q-$(A_2 + A_1)^{(1)}$ in subsection 2.

We use the following notations:

$$\begin{aligned} &a_1a_2\dots a_n/b_1b_2\dots b_n = \frac{a_1a_2\dots a_n}{b_1b_2\dots b_n},\\ &\mathcal{N}(x) = \prod_{i=0}^{m+n-1}(1-x/q^i),\\ &T_{a_i}(F) = F|_{a_i\to qa_i},\quad T_{a_i}^{-1}(F) = F|_{a_i\to a_i/q}, \end{aligned} \tag{3.1}$$

for any quantity (or function) F depending on variables a_i and b_i.

3.1. **Case** q-$E_7^{(1)}$**.**

(a) Setting of the Padé interpolation problem

The interpolated function, the interpolated sequence and the constraint (2.2):

$$Y(x) = \prod_{i=1}^{3}\frac{(a_ix,b_i;q)_\infty}{(a_i,b_ix;q)_\infty},\quad Y_s = \prod_{i=1}^{3}\frac{(b_i;q)_s}{(a_i;q)_s},\quad \frac{a_1a_2a_3q^m}{b_1b_2b_3q^n} = 1. \tag{3.2}$$

The time evolution (2.5):

$$T : (a_1,a_2,a_3,b_1,b_2,b_3,m,n)\mapsto(qa_1,a_2,qa_3,b_1,b_2,qb_3,m-1,n). \tag{3.3}$$

(b) Contiguity relations

The basic quantities:

$$G(x) = \prod_{i=1}^{3}\frac{(1-b_ix)}{(1-a_ix)},\quad K(x) = \frac{1-b_3x}{1-b_3}\prod_{i=1,3}\frac{(1-a_i)}{(1-a_ix)},\quad H(x) = (1-b_3)\prod_{i=1}^{3}(1-a_ix). \tag{3.4}$$

The Casorati determinants:

$$\begin{aligned} &D_1(x) = \frac{c_0x(1-xf)\mathcal{N}(x)Y(x)}{G_{\mathrm{den}}(x)},\quad D_2(x) = \frac{c_1(1-b_3x/a_2q^mg)\mathcal{N}(x)Y(x)}{K_{\mathrm{den}}(x)},\\ &D_3(x) = \frac{c_1(1-b_3x)(1-x/g)\mathcal{N}(x)Y(x)}{H(x)}, \end{aligned} \tag{3.5}$$

where f, g, c_0 and c_1 are constants with respect to x.

The contiguity relations:

$$
\begin{aligned}
&L_2(x) : C_0 x(1-xf)\overline{y}(x) - (1-a_2x)(1-b_3x/a_2q^m g)y(qx) + (1-b_3x)(1-x/g)y(x) = 0,\\
&L_3(x) : C_1 x(1-x\overline{f}/q)y(x) + \frac{A_2(x)}{1-b_3x}(1-\frac{x}{qg})\overline{y}(x) - \frac{A_1(x/q)}{1-a_2x/q}(1-b_3x/a_2q^m g)\overline{y}(x/q) = 0,
\end{aligned}
\tag{3.6}
$$

where

$$
\begin{aligned}
&A_1(x) = (1-a_2x)(1-qx)\prod_{i=1,2}(1-b_ix),\\
&A_2(x) = (1-b_3x)(1-x/q^{m+n})\prod_{i=1,3}(1-a_ix),\\
&C_0 = c_0(1-b_3)/c_1, \quad C_1 = \overline{c}_0(1-a_1)(1-a_3)/qc_1.
\end{aligned}
\tag{3.7}
$$

(c) The q-Painlevé equation

Compatibility gives the following equations:

$$
\begin{aligned}
&\frac{(fg-1)(f\underline{g}-1)}{(fg-b_3/a_2q^m)(f\underline{g}-b_3/a_2q^{m+1})} = \frac{A_1(1/f)}{A_2(1/f)},\\
&\frac{(1-fg)(1-\overline{f}g)}{(1-a_2q^m fg/b_3)(1-a_2q^{m-1}\overline{f}g/b_3)} = \frac{A_1(g)}{A_2(a_2q^m g/b_3)}.
\end{aligned}
\tag{3.8}
$$

These equations (3.8) are equivalent to the q-Painlevé equation of type $E_7^{(1)}$ given in [4] [9] [17]. The 8 singular points are on the two curves $fg = 1$ and $fg = b_3/a_2q^m$.

$$
\begin{aligned}
(f,g) = {}&(a_2, 1/a_2), (b_1, 1/b_1), (b_2, 1/b_2), (q, 1/q),\\
&(a_1, b_3/a_1a_2q^m), (b_3, 1/a_2q^m), (1/q^{m+n}, b_3q^n/a_2), (a_3, b_3/a_2a_3q^m).
\end{aligned}
\tag{3.9}
$$

The product C_0C_1:

$$
C_0C_1 = \frac{A_1(g)(1-b_3/a_2q^m)(1-b_3/a_2q^{m-1})}{q(1-fg)(1-\overline{f}g)g^2}.
\tag{3.10}
$$

(d) The Lax pair

$$
\begin{aligned}
L_1(x) : {}&\frac{(b_3-a_2q^m)x^2}{a_2b_3q^m g}\left[\frac{A_1(g)}{(fg-1)(qg-x)} - \frac{A_2(a_2q^m g/b_3)}{(a_2q^m fg/b_3-1)(a_2q^m g/b_3-x)}\right]y(x)\\
&+ \frac{q^2(q-b_3x)A_1(x/q)}{(q-a_2x)(q-fx)}\left[y(x/q) - \frac{b_3(a_2q^{m+1}g/b_3-x)(q-a_2x)}{a_2q^m(qg-x)(q-b_3x)}y(x)\right]\\
&+ \frac{(1-a_2x)A_2(x)}{(1-b_3x)(1-fx)}\left[y(qx) - \frac{a_2q^m(g-x)(1-b_3x)}{b_3(a_2q^m g/b_3-x)(1-a_2x)}y(x)\right] = 0,\\
L_2(x) : {}&C_0x(1-xf)\overline{y}(x) - (1-a_2x)(1-b_3x/a_2q^m g)y(qx)\\
&+ (1-b_3x)(1-x/g)y(x) = 0.
\end{aligned}
\tag{3.11}
$$

The scalar Lax pair (3.11) is equivalent to that in [23] using a suitable gauge transformation of $y(x)$. (Note that there is a typographical error in the second equation of (36) in [23], namely the expression $\overline{f}g - t^2$ should read $\overline{f}gq - t^2$.)

(e) Special solutions

$$\frac{1-f/a_1}{1-f/a_2}=\frac{\gamma_1 T_{a_1}(\tau_{m,n})T_{a_1}^{-1}(\tau_{m+1,n-1})}{\gamma_2 T_{a_2}(\tau_{m,n})T_{a_2}^{-1}(\tau_{m+1,n-1})},\quad \frac{1-1/b_1 g}{1-1/b_2 g}=\frac{\omega_1 T_{b_1}^{-1}(\tau_{m,n})T_{b_1}(\overline{\tau}_{m+1,n-1})}{\omega_2 T_{b_2}^{-1}(\tau_{m,n})T_{b_2}(\overline{\tau}_{m+1,n-1})}, \tag{3.12}$$

where

$$\tau_{m,n}=\det\Big[{}_4\varphi_3\Big(\begin{matrix}b_1,b_2,b_3,q^{-(m+n)}\\ a_1,a_2,a_3\end{matrix};q,q^{i+j+1}\Big)\Big]_{i,j=0}^{n}, \tag{3.13}$$

$$\gamma_i=\frac{a_i(1-a_iq^{m+n})(1-a_i/q)^n\prod_{k=1}^{3}(1-b_k/a_i)}{(1-a_i)^{n+1}},\quad \omega_i=\frac{(1-a_2/b_i)(1-b_i)^n}{(1-b_i/q)^n},\quad \text{for } i=1,2. \tag{3.14}$$

These determinant formulae of hypergeometric solutions (3.12) are expressed in terms of the terminating balanced ${}_4\varphi_3$ (2.4) series (Askey–Wilson polynomials [9]), and they are expected to be equivalent to the terminating case of that in [13].

3.2. **Case** q-$E_6^{(1)}$.

The contents of this subsubsection is the same as [6].

(a) Setting of the Padé interpolation problem

The interpolated function and the interpolated sequence (2.2):

$$Y(x)=\prod_{i=1}^{2}\frac{(a_ix,b_i;q)_\infty}{(a_i,b_ix;q)_\infty},\quad Y_s=\prod_{i=1}^{2}\frac{(b_i;q)_s}{(a_i;q)_s}. \tag{3.15}$$

The time evolution (2.5):

$$T:(a_1,a_2,b_1,b_2,m,n)\mapsto(qa_1,a_2,,b_1,b_2,m-1,n). \tag{3.16}$$

(b) Contiguity relations

The basic quantities:

$$G(x)=\prod_{i=1}^{2}\frac{(1-b_ix)}{(1-a_ix)},\quad K(x)=\frac{1-a_1}{1-a_1x},\quad H(x)=\prod_{i=1}^{2}(1-a_ix). \tag{3.17}$$

The Casorati determinants:

$$D_1(x)=\frac{c_0x(1-xf)\mathcal{N}(x)Y(x)}{G_{\mathrm{den}}(x)},\quad D_2(x)=\frac{c_1\mathcal{N}(x)Y(x)}{K_{\mathrm{den}}(x)},\quad D_3(x)=\frac{c_1(1-x/g)\mathcal{N}(x)Y(x)}{H(x)}, \tag{3.18}$$

where f,g,c_0 and c_1 are constants with respect to x.

The contiguity relations:

$$\begin{aligned}&L_2(x):C_0x(1-xf)\overline{y}(x)-(1-a_2x)y(qx)+(1-x/g)y(x)=0,\\ &L_3(x):C_1x(1-x\overline{f}/q)y(x)+(1-a_1x)(1-x/q^{m+n})(1-x/qg)\overline{y}(x)\\ &\qquad\qquad-(1-x)(1-b_1x/q)(1-b_2x/q)\overline{y}(x/q)=0,\end{aligned} \tag{3.19}$$

where $C_0=c_0/c_1$ and $C_1=\overline{c}_0(1-a_1)/qc_1$.

(c) The q-Painlevé equation

Compatibility gives the following equations:

$$\begin{aligned}\frac{(fg-1)(f\underline{g}-1)}{g\underline{g}} &= \frac{(f-a_2)(f-b_1)(f-b_2)(f-q)}{(f-a_1)(f-1/q^{m+n})},\\ \frac{(fg-1)(\overline{f}g-1)}{f\overline{f}} &= \frac{(g-1/a_2)(g-1/b_1)(g-1/b_2)(g-1/q)}{(g-1/a_2q^m)(g-a_1/b_1b_2q^n)}.\end{aligned} \tag{3.20}$$

These equations (3.20) are equivalent to the q-Painlevé equation of type $E_6^{(1)}$ given in [6] [9] [17]. The 8 singular points are on the two lines $f=0$ and $g=0$ and one curve $fg=1$.

$$\begin{aligned}(f,g) = (a_2,1/a_2),(b_1,1/b_1),(b_2,1/b_2),(q,1/q),\\ (a_1,0),(1/q^{m+n},0),(0,1/a_2q^m),(0,a_1/b_1b_2q^n).\end{aligned} \tag{3.21}$$

The product C_0C_1:

$$C_0C_1 = \frac{(1-a_2g)(1-b_1g)(1-b_2g)(1-qg)}{qg^2(1-fg)(1-\overline{f}g)}. \tag{3.22}$$

(d) The Lax pair

(3.23)

$$\begin{aligned}L_1(x):\ &\frac{q^{m+n}g(1-x)(q-b_1x)(q-b_2x)}{q-fx}\left[y(x/q)-\frac{g(q-a_2x)}{qg-x}y(x)\right]\\ &+\frac{g(q^{m+n}-x)(1-a_1x)(1-a_2x)}{1-fx}\left[y(qx)-\frac{(g-x)}{g(1-a_2x)}y(x)\right]\\ &+x^2\left[\frac{(a_2q^mg-1)(b_1b_2q^ng-a_1)}{f}-\frac{q^{m+n}(a_2g-1)(b_1g-1)(b_2g-1)(qg-1)}{(fg-1)(qg-x)}\right]y(x)=0,\\ L_2(x):\ &C_0x(1-xf)\overline{y}(x)-(1-a_2x)y(qx)+(1-x/g)y(x)=0.\end{aligned}$$

The scalar Lax pair (3.23) is equivalent to the 2×2 matrix ones in [21] [27] and the scalar ones in [6] [23] using suitable gauge transformations of $y(x)$. (Note that there are some typographical errors in equations (30) and (31) in [6], namely the expressions $(b_4q'-z)$ and $Y(x)$ should read $(b_4q'-z)t^2$ and $Y(z)$ respectively.)

(e) Special solutions

$$\frac{1-f/a_1}{1-f/a_2}=\frac{\gamma_1T_{a_1}(\tau_{m,n})T_{a_1}^{-1}(\tau_{m+1,n-1})}{\gamma_2T_{a_2}(\tau_{m,n})T_{a_2}^{-1}(\tau_{m+1,n-1})},\quad \frac{1-1/b_1g}{1-1/b_2g}=\frac{\omega_1T_{b_1}^{-1}(\tau_{m,n})T_{b_1}(\overline{\tau}_{m+1,n-1})}{\omega_2T_{b_2}^{-1}(\tau_{m,n})T_{b_2}(\overline{\tau}_{m+1,n-1})}, \tag{3.24}$$

where

$$\tau_{m,n}=\det\left[{}_3\varphi_2\left(\begin{matrix}b_1,b_2,q^{-(m+n)}\\ a_1,a_2\end{matrix};q,q^{i+j+1}\right)\right]_{i,j=0}^{n}, \tag{3.25}$$

$$\gamma_i=\frac{a_i(1-a_iq^{m+n})(1-a_i/q)^n\prod_{k=1}^{2}(1-b_k/a_i)}{(1-a_i)^{n+1}},\quad \omega_i=\frac{(1-a_2/b_i)(1-b_i)^n}{(1-b_i/q)^n},\quad \text{for } i=1,2. \tag{3.26}$$

These determinant formulae of hypergeometric solutions (3.24) are expressed in terms of the terminating ${}_3\varphi_2$ (2.4) series (big q-Jacobi polynomials [9]), and they are equivalent to that in [6]. (Note that there is a typographical error in equation (38) in [6], namely the expression $T_{a_2}T_{a_3}(\tau_{m,n-1})$ should read $T_{a_2}T_{a_4}(\tau_{m,n-1})$.)

3.3. **Case q-$D_5^{(1)}$.**

(a) Setting of the Padé interpolation problem

The interpolated function and the interpolated sequence (2.2):

$$Y(x) = c^{\log_q x}\frac{(a_1x, b_1; q)_\infty}{(a_1, b_1x; q)_\infty}, \quad Y_s = c^s\frac{(b_1; q)_s}{(a_1; q)_s}. \tag{3.27}$$

The time evolution (2.5):

$$T : (a_1, b_1, c, m, n) \mapsto (qa_1, b_1, c, m-1, n). \tag{3.28}$$

(b) Contiguity relations

The basic quantities:

$$G(x) = \frac{(1-b_1x)c}{1-a_1x}, \quad K(x) = \frac{1-a_1}{1-a_1x}, \quad H(x) = 1-a_1x. \tag{3.29}$$

The Casorati determinants:

$$D_1(x) = \frac{c_0(1-xf)\mathcal{N}(x)Y(x)}{G_{\mathrm{den}}(x)}, \quad D_2(x) = \frac{c_1N(x)Y(x)}{K_{\mathrm{den}}(x)}, \quad D_3(x) = \frac{c_2N(x)Y(x)}{H(x)}, \tag{3.30}$$

where f, c_0, c_1 and c_2 are constants with respect to x.
The contiguity relations:

$$\begin{aligned} &L_2(x) : C_0(1-xf)\overline{y}(x) - y(qx) + y(x)/g = 0, \\ &L_3(x) : C_1(1-x\overline{f}/q)y(x) - (1-a_1x)(1-x/q^{m+n})\overline{y}(x)/g \\ &\qquad\qquad + c(1-x)(1-b_1x/q)\overline{y}(x/q) = 0, \end{aligned} \tag{3.31}$$

where $C_0 = c_0/c_1, C_1 = -\overline{c}_0(1-a_1)/c_1$ and $g = c_1/c_2$.

(c) The q-Painlevé equation

Compatibility gives the following equations:

$$g\underline{g} = \frac{(f-a_1)(f-1/q^{m+n})}{c(f-b_1)(f-q)}, \quad f\overline{f} = \frac{qb_1(g-1/q^m)(g-a_1/b_1q^nc)}{(g-1)(g-1/c)}. \tag{3.32}$$

These equations (3.32) are equivalent to the q-Painlevé equation of type $D_5^{(1)}$ given in [8] [9]. The 8 singular points are on the four lines $f = 0$, $f = \infty$, $g = 0$ and $g = \infty$.

$$(f, g) = (a_1, 0), (1/q^{m+n}, 0), (0, 1/q^m), (0, a_1/b_1cq^n), (\infty, 1), (\infty, 1/c), (b_1, \infty), (q, \infty). \tag{3.33}$$

The product C_0C_1:

$$C_0C_1 = -(1-g)(1-cg)/g^2. \tag{3.34}$$

(d) The Lax pair

$$\begin{aligned} L_1(x) : &\frac{g(x-q^{m+n})(a_1x-1)}{fx-1}\left[y(qx) - \frac{y(x)}{g}\right] + \frac{cq^{m+n}g(x-1)(b_1x-q)}{fx-q}\left[y(\frac{x}{q}) - gy(x)\right] \\ &+ \left[\frac{(q^mg-1)(b_1cq^ng-a_1)x}{f} - q^{m+n}(g-1)(cg-1)\right]y(x) = 0, \\ L_2(x) : &C_0(1-xf)\overline{y}(x) - y(qx) + y(x)/g = 0. \end{aligned} \tag{3.35}$$

The scalar Lax pair (3.35) is equivalent to the 2×2 matrix ones in [8] [14] and the scalar one in [23] using suitable gauge transformations of $y(x)$.

(e) Special solutions

$$\begin{aligned} \frac{f}{a_1} - 1 &= \frac{c(1-b_1/a_1)(1-a_1q^{m+n})(1-a_1/q)^n}{q^m(1-c)(1-a_1)^{n+1}} \frac{T_{a_1}(\tau_{m,n,1})T_{a_1}^{-1}(\tau_{m+1,n-1,1})}{\tau_{m,n,0}\tau_{m+1,n-1,2}}, \\ g &= \frac{(1-a_1q^{m+n})(1-b_1/q)^n}{q^m(1-a_1)(1-b_1)^n} \frac{T_{a_1}(\tau_{m,n,1})\tau_{m,n-1,1}}{T_{b_1}^{-1}(\tau_{m,n,1})T_{a_1b_1}(\tau_{m,n-1,1})}, \end{aligned} \tag{3.36}$$

where

$$\tau_{m,n,k} = \det\left[{}_2\varphi_1\left(\begin{matrix} b_1, q^{-(m+n)} \\ a_1 \end{matrix}; q, cq^{i+j+k}\right)\right]_{i,j=0}^{n}. \tag{3.37}$$

These determinant formulae of hypergeometric solutions (3.36) are expressed in terms of the terminating ${}_2\varphi_1$ (2.4) series (little q-Jacobi polynomials [9]), and they are expected to be equivalent to the terminating case of that in [18].

3.4. **Case** q-$A_4^{(1)}$**.**

(a) Setting of the Padé interpolation problem

The interpolated function and the interpolated sequence (2.2):

$$Y(x) = c^{\log_q x}\frac{(b_1;q)_\infty}{(b_1x;q)_\infty}, \quad Y_s = c^s(b_1;q)_s. \tag{3.38}$$

The time evolution (2.5):

$$T : (b_1, c, m, n) \mapsto (b_1, c, m-1, n). \tag{3.39}$$

(b) Contiguity relations

The basic quantities:

$$G(x) = (1-b_1x)c, \quad K(x) = 1, \quad H(x) = 1. \tag{3.40}$$

The Casorati determinants:

$$D_1(x) = \frac{c_0(1-xf)\mathcal{N}(x)Y(x)}{G_{\mathrm{den}}(x)}, \quad D_2(x) = \frac{c_1N(x)Y(x)}{K_{\mathrm{den}}(x)}, \quad D_3(x) = \frac{c_2N(x)Y(x)}{H(x)}, \tag{3.41}$$

where f, c_0, c_1 and c_2 are constants with respect to x.

The contiguity relations:

$$\begin{aligned} &L_2(x) : C_0(1-xf)\overline{y}(x) - y(qx) + y(x)/g = 0, \\ &L_3(x) : C_1(1-x\overline{f}/q)y(x) - (1-x/q^{m+n})\overline{y}(x)/g + c(1-x)(1-b_1x/q)\overline{y}(x/q) = 0, \end{aligned} \tag{3.42}$$

where $C_0 = c_0/c_1$, $C_1 = -\overline{c}_0/c_1$ and $g = c_1/c_2$.

(c) The q-Painlevé equation

Compatibility gives the following equations:

$$g\underline{g} = \frac{f(f-1/q^{m+n})}{c(f-b_1)(f-q)}, \quad f\overline{f} = \frac{qb_1g(g-1/q^m)}{(g-1)(g-1/c)}. \tag{3.43}$$

These equations (3.43) are equivalent to the q-Painlevé equation of type $A_4^{(1)}$ given in [9] [11]. The 8 singular points are on the four lines $f = 0$, $f = \infty, g = 0$ and $g = \infty$. $(0, 0)$ is a double point.

$$\begin{aligned}(f, g) &= (1/q^{m+n}, 0), (0, 1/q^m), (\infty, 1), (\infty, 1/c), (b_1, \infty), (q, \infty),\\ (0, 0) &\leftarrow g/f = -1/b_1 c q^n.\end{aligned} \tag{3.44}$$

The product C_0C_1:

$$C_0C_1 = -(1 - g)(1 - cg)/g^2. \tag{3.45}$$

(d) The Lax pair

$$\begin{aligned}L_1(x) :\ & \frac{cgq^{m+n}(1 - x)(q - b_1x)}{q - fx}\left[y(x/q) - gy(x)\right] + \frac{g(q^{m+n} - x)}{1 - fx}\left[y(qx) - y(x)/g\right]\\ &+ q^n\left[q^m(1 - g)(1 - cg) + b_1cg(1 - q^m g)x/f\right]y(x) = 0,\\ L_2(x) :\ & C_0(1 - xf)\overline{y}(x) - y(qx) + y(x)/g = 0.\end{aligned} \tag{3.46}$$

The scalar Lax pair (3.46) is equivalent to the 2×2 matrix one for the q-Painlevé equation of type q-$P(A_4)$ in [14] using a suitable gauge transformation of $y(x)$.

(e) Computation of special solutions

$$f = \frac{b_1c}{(c - 1)q^m}\frac{\tau_{m,n,1}\tau_{m+1,n-1,1}}{\tau_{m,n,0}\tau_{m+1,n-1,2}}, \quad g = \frac{(1 - b_1/q)^n}{q^m(1 - b_1)^n}\frac{\tau_{m,n,1}\tau_{m,n-1,1}}{T_{b_1}^{-1}(\tau_{m,n,1})T_{b_1}(\tau_{m,n-1,1})}, \tag{3.47}$$

where

$$\tau_{m,n,k} = \det\left[{}_2\varphi_1\left(\begin{matrix}b_1, q^{-(m+n)}\\ 0\end{matrix}; q, cq^{i+j+k}\right)\right]_{i,j=0}^{n}. \tag{3.48}$$

These determinant formulae of hypergeometric solutions (3.47) are expressed in terms of the terminating ${}_2\varphi_1$ (2.4) series (q-Laguerre polynomials [9]), and they are expected to be equivalent to the terminating case of that in [5].

3.5. **Case** q-$(A_2 + A_1)^{(1)}$.

(a) Setting of the Padé interpolation problem

The interpolated function and the interpolated sequence (2.2):

$$Y(x) = (d\sqrt{x/q})^{\log_q x}, \quad Y_s = q^{\binom{s}{2}}d^s. \tag{3.49}$$

The time evolution (2.5):

$$T : (d, m, n) \mapsto (d, m - 1, n). \tag{3.50}$$

(b) Contiguity relations

The basic quantities:

$$G(x) = dx, \quad K(x) = 1, \quad H(x) = 1. \tag{3.51}$$

The Casorati determinants:

$$D_1(x) = \frac{c_0(1 - xf)\mathcal{N}(x)Y(x)}{G_{\mathrm{den}}(x)}, \quad D_2(x) = \frac{c_1N(x)Y(x)}{K_{\mathrm{den}}(x)}, \quad D_3(x) = \frac{c_2N(x)Y(x)}{H(x)}, \tag{3.52}$$

where f, c_0, c_1 and c_2 are constants with respect to x.

The contiguity relations:

$$\begin{aligned} &L_2(x) : C_0(1 - xf)\overline{y}(x) - y(qx) + y(x)/g = 0, \\ &L_3(x) : C_1(1 - x\overline{f}/q)y(x) + (1 - x/q^{m+n})\overline{y}(x)/g - dx(1 - x)\overline{y}(x/q)/q = 0, \end{aligned} \tag{3.53}$$

where $C_0 = c_0/c_1$, $C_1 = \overline{c}_0/c_1$ and $g = c_1/c_2$.

(c) The q-Painlevé equation

Compatibility gives the following equation:

$$g\underline{g} = \frac{f(f - 1/q^{m+n})}{d(f - q)}, \quad f\overline{f} = \frac{qdg(g - 1/q^m)}{(g - 1)}. \tag{3.54}$$

These equations (3.54) are equivalent to the q-Painlevé equation of type $(A_2 + A_1)^{(1)}$, namely P_{III}, given in [9] [11]. The 8 singular points are on the four lines $f = 0, f = \infty, g = 0$ and $g = \infty$. The two points $(0, 0)$ and (∞, ∞) are double points.

$$\begin{aligned} (f, g) &= (1/q^{m+n}, 0), (0, 1/q^m), (\infty, 1), (q, \infty), \\ &(0, 0) \leftarrow g/f = 1/dq^n, \\ &(\infty, \infty) \leftarrow g/f = 1/d. \end{aligned} \tag{3.55}$$

The product C_0C_1:

$$C_0C_1 = (1 - g)/g^2. \tag{3.56}$$

(d) The Lax pair

$$\begin{aligned} L_1(x) : &\frac{dgq^{m+n}(1 - x)x}{fx - q}\left[y(x/q) - gy(x)\right] + \frac{g(q^{m+n} - x)}{fx - 1}\left[y(qx) - y(x)/g\right] \\ &+ q^n\left[q^m(g - 1) - dg(gq^m - 1)x/f\right]y(x) = 0, \\ L_2(x) : &C_0(1 - xf)\overline{y}(x) - y(qx) + y(x)/g = 0. \end{aligned} \tag{3.57}$$

The scalar Lax pair (3.57) is equivalent to the 2×2 matrix one for the q-Painlevé equation of type q-$P(A_5)^{\#}$ in [14] using a suitable gauge transformation of $y(x)$.

(e) Special solutions

$$f = \frac{d}{q^m}\frac{\tau_{m,n,1}\tau_{m+1,n-1,1}}{\tau_{m,n,0}\tau_{m+1,n-1,2}}, \quad g = \frac{1}{q^{m+n}}\frac{\tau_{m,n,1}\tau_{m,n-1,1}}{\tau_{m,n,0}\tau_{m,n-1,2}}, \tag{3.58}$$

where

$$\tau_{m,n,k} = \det\left[{}_1\varphi_1\left(\begin{matrix} q^{-(m+n)} \\ 0 \end{matrix}; q, -dq^{i+j+k}\right)\right]_{i,j=0}^{n}. \tag{3.59}$$

These determinant formulae of hypergeometric solutions (3.58) are expressed in terms of the terminating ${}_1\varphi_1$ (2.4) series (Stieltjes–Wigert polynomials [9]), and they are expected to be equivalent to the terminating case of that in [15].

4. Conclusion

4.1. **Summary.**

In this paper, for the *interpolated function* $Y(x)$ given in Table (2.2), we established the *Padé interpolation problem* related to all the q-Painlevé equations of types from $E_7^{(1)}$ to $(A_2 + A_1)^{(1)}$. Then, for the *time evolution* T given in Table (2.5), we established another Padé interpolation problem. By solving these problems, we derived the evolution equations, the scalar Lax pairs and the determinant formulae of the special solutions for the corresponding q-Painlevé equations. The main results are given in section 3.

4.2. **Problems.**

Some open problems related to the results of this paper are as follows:

1. In this paper, by choosing one time evolution T, we applied the Padé method to each type from q-$E_7^{(1)}$ to q-$(A_2 + A_1)^{(1)}$. By choosing other time evolutions, we can perform similar computations. It may be interesting to study the relation between the Padé method for the various time evolutions and Bačklund transformations of the affine Weyl group, for example the case q-$E_6^{(1)}$ in [6].

2. By the results of this paper and previous ones [6][16][24], it turned out that the Padé method could be applied to all the discrete Painlevé equations of types from e-$E_8^{(1)}$ to q-$(A_2 + A_1)^{(1)}$. It may be interesting to study the degenerations between these results.

3. In this paper, we applied the Padé method of the q-grid to the q-Painlevé equation of type $D_5^{(1)}$. On the other hand, the method of the differential grid (i.e., Padé approximation) was also applied to the q-P_{VI} equation in [6][25]. It may be interesting to study whether the method of the differential grid can be also applied to the q-Painlevé equations of other types.

4. It may be interesting to study whether the Padé method can be further applied to the other generalized Painlevé systems, for example the q-Garnier system in [20].

Acknowledgment

The author is grateful to Professor Y. Yamada for valuable discussions on this research, and for his encouragement. The author also thanks Professors T. Suzuki, T. Tsuda, H. Sakai, M. Noumi, W. Rossman and the referee for stimulating comments and for kindhearted support.

References

[1] Clarkson Peter A., *Recurrence coefficients for discrete orthonormal polynomials and the Painlevé equations*, J. Phys., A **46**, no.18, (2013), 185205(18pp).

[2] Gasper G. and Rahman M., *Basic Hypergeometric Series*, Encyclopedia of Mathematics and Its Applications, Volume **35** Cambridge University Press, Cambridge (1990)

[3] Grammaticos B., Nijhoff F.W., Ramani A., *Discrete Painlevé equations, The Painlevé Property: One Century Later*, Editor R. Conte, CRM Ser. Math. Phys., Springer, New York, (1999), 413–516.

[4] Grammaticos B., Ramani A., *On a novel q-discrete analogue of the Painlevé* VI equation, Phys. Lett., A**257** (1999), 288–292.

[5] Hamamoto T., and Kajiwara K., *Hypergeometric solutions to the q-Painlevé equation of type* $A_4^{(1)}$, J. Phys. A: Math. Theor., **40** (2007), 12509–12524.

[6] Ikawa Y., *Hypergeometric Solutions for the q-Painlevé Equation of Type* $E_6^{(1)}$ *by the Padé method*, Lett. Math. Phys., Volume **103**, Issue 7 (2013), 743–763.

[7] Jacobi,C.G.J Über die Darstellung einer Reihe gegebner Werthe durch eine gebrochne rationale Function. J. Reine Angew. Math., **30** (1846) 127–156.

[8] Jimbo M. and Sakai H., *A q-analog of the sixth Painlevé equation*, Lett. Math. Phys., **38** (1996), 145–154.

[9] Kajiwara K., Masuda T, Noumi M., Ohta Y. and Yamada Y., *Hypergeometric solutions to the q-Painlevé equations*, IMRN 2004, **47** (2004), 2497-2521.

[10] Kajiwara K., Noumi M., and Yamada Y., *A study on the fourth q-Painlevé equation*, J. Phys. A: Math. Gen., **34** (2001), 8563–8581.

[11] Kruskal M. D., Tamizhmani K. M., B. Grammaticos and Ramani A., *Asymmetric discrete Painlevé equations*, Regul. Chaot. Dyn., **5** (2000), 273–281.
[12] Magnus A., *Painlevé-type differential equations for the recurrence coefficients of semi- classical orthogonal polynomials*, J. Comput. Appl. Math., **57** (1995), 215-237.
[13] Masuda T., *Hypergeometric τ-functions of the q-Painlevé system of type $E_7^{(1)}$*, SIGMA, **5** (2009), 035 (30pp).
[14] Murata M., *Lax forms of the q-Painlevé equations*, J. Phys. A: Math. Theor., **42** (2009), 115201(17pp.).
[15] Nakazono N., *Hypergeometric tau Functions of the q-Painlevé Systems of Type $(A_2+A_1)^{(1)}$* SIGMA, **6** (2010), 084 (16pp).
[16] Noumi M., Tsujimoto S., Yamada Y., *Padé interpolation for elliptic Painlevé equation*, Symmetries, integrable systems and representations, Springer Proc. Math. Stat., Volume **40** (2013), 463–482.
[17] Ramani A., Grammaticos B., Tamizhmani T., and Tamizhmani K.M., Special Function Solutions of the Discrete Painlevé Equations, Comput. Math. Appl., **42** (2001), no. 3–5, 603–614.
[18] Sakai H., *Casorati determinant solutions for the q-difference sixth Painlevé equations*, Nonlinearity, **11** (1998), 823–833.
[19] Sakai H., *Rational surfaces with affine root systems and geometry of the Painlevé equations*, Commun. Math. Phys., **220** (2001), 165–221.
[20] Sakai H., *A q-Analog of the Garnier System*, Funkcial. Ekvac. **48** (2005), 273–297.
[21] Sakai H., *Lax form of the q-Painlevé equation associated with the $A_2^{(1)}$ surface*, J. Phys. A: Math. Gen., **39** (2006), 12203-12210.
[22] Yamada Y., *Padé method to Painlevé equations*, Funkcial. Ekvac., **52** (2009), 83–92.
[23] Yamada Y., *Lax formalism for q-Painlevé equations with affine Weyl group symmetry of type $E_n^{(1)}$*, IMRN 2011, **17** (2011), 3823–3838.
[24] Yamada Y., *A simple expression for discrete Painlevé equations*, RIMS Kokyuroku Bessatsu, **B47** (2014), 087–095, to appear.
[25] Yoshioka R., *Padé approximation and special solution for q-Painlevé VI equation*. Master thesis in Kobe University (Japanese) (2010)
[26] Van Assche.W., *Discrete Painlevé equations for recurrence coefficients of orthogonal polynomials. Difference equations, special functions and orthogonal polynomials*, World Sci. Publ., Hackensack, NJ (2007) 687–725.
[27] Witte N.S, Ormerod C.M, *Construction of a Lax Pair for the $E_6^{(1)}$ q-Painlevé System*, SIGMA, **8** (2012), 097 (27pp).
[28] Zhedanov A.S., *Padé interpolation table and biorthogonal rational functions*, Rokko Lectures in Mathematics, **18** (2005), 323–363.

Department of Mathematics, Graduate School of Science, Kobe University, Hyogo 657-8501, Japan
E-mail address: hnagao@math.kobe-u.ac.jp